



A Comprehensive Review on Cytoskeletal Organization and Its Interaction with Polysaccharides and Enzymes during Primary Cell Wall Development

Rao Saad Rehman ^{a*}, Hassan Bashir ^b, Syed Ali Zafar ^c, Mujahid Ali ^d,
Abdullah Javed ^e, Naveed Ali Ashraf ^e, Muzammal Hussain ^f
and Ayesha Hussain ^f

^a College of Plant Science and Technology, Huazhong Agricultural University, Wuhan, Hubei, China.

^b Department of Plant Breeding and Genetics, Sub Campus Burewala-Vehari, University of Agriculture Faisalabad, Pakistan.

^c Oilseeds Research Institute, Ayub Agricultural Research Institute, Faisalabad, Pakistan.

^d Department of Plant Breeding and Genetics, Nanjing Agricultural University, Nanjing, China.

^e Department of Seed Science and Technology, University of Agriculture Faisalabad, Pakistan.

^f Department of Soil and Environmental Sciences, Sub Campus Burewala-Vehari, University of Agriculture Faisalabad, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BJI/2022/v26i4653

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91895>

Review Article

Received 17 July 2022
Accepted 23 September 2022
Published 26 September 2022

ABSTRACT

A well-coordinated process is required to construct a complicated structure like the cell wall, which consists of several elements that must be joined appropriately from various sources inside the cell. In order to successfully moderate dynamic responses to developmental and environmental signals, further complexity is necessary. The plasma membrane is continually and actively transporting sugars, enzymes, and other cell wall elements throughout diffused development. Actin filaments and microtubules make up the cytoskeletal pathways used to transport cell wall elements in vesicles during cell division. In addition to these elements, other proteins, vesicles and lipids are transported

*Corresponding author: E-mail: saad_rehman101@hotmail.com;

from and to the cell plate while cytokinesis occurs. Adding additional cell wall material or building a new cell wall requires a rearrangement of the cytoskeleton, which we examine in this review first. We next look at the commonalities between these two processes. Our next topic is the transport of cell wall-building polysaccharides and enzymes via motor proteins and other interactions with the cytoskeleton. Final thoughts on cytokinesis-generated cell walls include a look at some of their unique properties.

Keywords: Cell cortex; cytoskeleton; phragmoplast; microtubule nucleators; cellulose biosynthesis; kinesins; callose.

1. INTRODUCTION

In 1665, when Robert Hooke used a microscope to investigate oak cork (*Quercus suber*) under a light microscope, he accidentally started the study of cell wall function and structure. Because of its resemblance to a monk's room (cell), he came up with the name "cell" to represent the tiniest living biological structure. Hooke was able to see the cell wall for the first time since the cellular contents had been cleared. Cell wall research has since profited from structural component analysis of the isolated cell wall [1]. Our knowledge of how the cytoskeleton and transport of cell wall components are intertwined has recently improved thanks to recent advances in cell biology, especially live-cell imaging (LCI). Cell expansion and division are two of the most important factors in generating plant biomass. Cellulose is the most prevalent biopolymer on the earth because it is a key element of main cell walls. Additionally, cell wall synthesis and alteration are as important to plants as global biomass. Cell wall production and modification take up around 10% of certain plant genomes, and hundreds to thousands of additional genes are involved in controlling the vesicle and cytoskeleton transportation [2].

During cytokinesis, the cell wall is built from scratch, whereas during diffuse growth, new cell wall material is incorporated into the existing cell wall. In both circumstances, motor proteins transport vesicles carrying cell wall components to the cytoskeleton. Vesicles normally emerge in the Golgi and are transported via the cytoskeleton, often passing through the trans Golgi network (TGN), on their way from and to the cell plate and the plasma membrane. For this reason, having microtubules and microfilaments that are appropriately structured is essential for the placement and assembly of cell wall substituents. For example, the quick and precise addition of cellulose during diffuse development is dependent on well-organized microtubules owing to direct connections between enzymes,

microtubule linking proteins and microtubules that create cell wall polymers called cellulose synthases.

During cell division, the phragmoplast, a conserved land-plant-specific structure, initiates the formation of the de novo cell wall [3]. Phragmoplasts are primarily responsible for generating new cell walls after the cytokinesis process separates the DNA [4]. Prior to the conclusion of cytokinesis, a preliminary version of the new cell wall is produced, known as the cell plate. As the cell plate grows in size, vesicle transport and fusion and changes in cell wall structure occur simultaneously [1,2,5]. Actin filaments in the phragmoplast create an antiparallel array of microtubules facing the center of the cell [6]. In the phragmoplast, spindle microtubules are used to build microtubules, which are then used to expand the phragmoplast through microtubule-dependent microtubule nucleation [7]. After then, when the cell plate grows, the microtubules are disassembled [6,8]. The cell plate fusion site or division site is a specified area where the phragmoplast connects to the cortex of the cell [3,9]. Because of this, the next section focuses on proteins and microfilaments that influence cytoskeletal dynamics and structure.

2. REGULATORY FACTORS RESPONSIBLE FOR MICROFILAMENT AND MICROTUBULAR BIOSYNTHESIS DURING DIFFUSE GROWTH

Actin filaments and microtubules are a pair in microfilaments, cargo, including cell wall components, are transported by motor proteins along the tracks. When the cell wall is changed, enlarged, or created, the factors that influence microfilament and microtubule placement have a significant but sometimes indirect function. The concept that cell shape controls orientation of microtubule is supported by the observation of cortical microtubules that are aligned with the long axis of rectangular or elliptical

microchambers [10]. In protoplasts, which are unicellular with enzymatically destroyed cell walls, the arrangement of microtubules was recently scientifically examined. Microtubules align along the long axis when protoplasts are restricted in rectangular wells, which suggests that the default orientation of microtubules is longitudinal [11]. *Planta*, on the other hand, often has microtubules that run perpendicular to the cell's long axis [12]. In order to transform the orientation of microtubules from longitudinal to transverse *in silico*, a faint directional signal is all that is required [13]. One of the directional signals is mechanical stress, which has been postulated. Cellulose microfibril orientation is guided by microtubules in response to mechanical stress, according to the most basic paradigm [14]. A cell growth path is perpendicular to the expected maximum tensile stress direction, whereas microtubules align parallel to this [12,15]. Microtubules develop circumferential in hypocotyls, cotyledons, and meristems when the stress pattern is altered by laser ablation of adjacent cells [16]. Spherical protoplasts were contained in rectangular reservoirs in solutions of varying osmolarity to further investigate the mechanism of microtubule orientation. Osmolarity decrease causes stress, which overrides geometrical signals that would otherwise direct tubular orientation along the projected path of maximum tension after two hours [7,17]. Unlike animal cells, plant cells are unlikely to experience comparable osmotic shifts, and it is not yet clear how the change in osmotic potential affects tensile stress. Even if microtubules are able to directly react to changes in tensile stress, numerous additional proteins are very certainly necessary. To find out whether regulators, the microtubules itself, or other components are involved in the detection of maximum tension, further study is required.

Formins have also had their roles clarified through a number of useful ways that target groupings of formins. In moss, RNAi suppressed all of the class 1 formins without affecting tip development, but plants sizes were shorter and had fewer cells, indicating delayed cytokinesis. Multinucleate or stubs cells were not found [18]. Understanding how formins contribute to cytokinesis and mitosis was made possible with the use of the small chemical formin inhibitor SMIFH2 [4-7,19]. Formins may occasionally be found in the phragmoplast midline or phragmoplast, despite their generally obscure

involvement in cytokinesis. This treatment also lowers duration of EB1 comet and changes turnover of fusion proteins such as KNOLLE and DYNAMIN RELATED PROTEIN 1A (DRP1A) [20]. No one knows for sure yet whether SMIFH2-induced cytokinesis deficits are primarily due to formin disruption. Formin mutations may be combined to offer clear evidence concerning their involvement in cell proliferation and cytokinesis.

3. REGULATORY FACTORS RESPONSIBLE FOR MICROFILAMENT AND MICROTUBULAR ORGANIZATION AT THE CELLULAR CORTEX DURING TELOPHASE

Cell cortex microtubules and actin filaments accumulate during telophase, but before the phragmoplast touches the cell cortex (Fig. 1) [21]. When microtubules emerge, they originate from either the nuclear envelope or the cortex-localized gamma tubulin or gamma tubulin ring complex [13,22]. A recent study has shown that the cell cortex may be organized by interactions with proteins that localize to the division site before the phragmoplast reaches the cortex, but it's not yet clear what these proteins do at this stage.

Through transitory microtubule plus-end stabilization, cortical telophase microtubules interact with TANGLED1 (TAN1) and other division site-localized proteins and this leads to the microtubules adopting a perpendicular orientation, which is analogous to the orientation of the phragmoplast itself [23]. Additionally, the *in vitro* involvement of TAN1 angle-independent crosslinking and bundling of microtubules is consistent with microtubule plus-end capture [18,24]. As the phragmoplast reaches the cell cortex, it incorporates microtubules from the cortical telophase. Parallel bundling of microtubules solely on a single part of the phragmoplast causes expansion of the phragmoplast to migrate toward the microtubules, which are subsequently incorporated into the phragmoplast via parallel bundling. There are yet no clues as to how additional microtubule-binding proteins that are present at the division site other than TAN1 engage with cortical telophase microtubules to fine tune the location of the phragmoplast to the division site.




Fig. 1. The cell cortex underlying the plasma membrane before it comes into contact with the phragmoplast

4. LOCALIZATION OF THE REGULATORY PROTEINS PRIOR TO PHRAGMOPLAST-CORTEX INTERACTION

Pre-cytokinesis proteins, those that concentrate at or near the division site, play an important function in cytokinesis and phragmoplast placement. Microtubule-cross-linking protein TAN1, kinesin PHRAGMOPLAST ORIENTING (POK1) and POK2 as well as IQ67 Domain (IQD) proteins are among the many other proteins that have been identified as having a role in mitosis. In maize, the protein TAN1 is necessary for proper placement of the phragmoplast, and it also bundles microtubules in vitro [25]. The AUXIN INDUCED IN ROOT CULTURES9 (AIR9) mutant, by itself, does not exhibit a phenotype that differs from that of the wild type in Arabidopsis tan1 mutants [26]. Double tan1 air9 mutants, on the other hand, have abnormalities in the division plane and have short roots [27]. The tan1 air9 double mutant wild-type development is restored by mitotic expression of TAN1 [28]. After the phragmoplast touches the cortex, the AIR9 protein concentrates at the division site [29]. After metaphase, two PH-GAPs that engage with POK1 are found to assemble at

the division site. These PH-GAPs are essential for situating the division plane in a redundant manner [30]. At last, a group of proteins known as "IQ67-DOMAIN" (IQD6, 7, and 8) interact with POK1 and have a role in establishing the division plane [31]. By examining where the proteins that constitute the division site are located in different mutants, we may learn more about how the division site is established and maintained until cytokinesis is complete.

5. CYTOKINESIS AND PHRAGMOPLAST CORTEX INTERACTIONS

Associations between microtubules and phragmoplast and actin filaments that are positioned in the cortex direct the phragmoplast exactly to the spot where it will divide after it has reached the cortex. They are facilitated by proteins located at division site, such as MYOSIN VIII in moss (microtubule and microfilament binding protein) [32]. Final contact with the cell cortex is dependent on the contacts between F-actin, Myosin, and the cell cortex. If these relationships are disrupted in any way, the phragmoplast will not be able to reach the mother cell plasma membrane [27,33]. Final contact with the cell cortex is dependent on the

contacts between F-actin, Myosin, and the cell cortex. If these relationships are disrupted in any way, the phragmoplast will not be able to reach the mother cell plasma membrane. The subsequent recruitment of the phragmoplast back to the division site suggests long distance interactions between the actin division sites [34]. In order for cells to properly extend and undergo cytokinesis, microtubule regulators including as nucleators, cross-linkers, end-binding proteins, and severing proteins are required [35].

6. REGULATORY ROLE OF MICROTUBULE NUCLEATORS DURING DIFFUSE GROWTH AND CYTOKINESIS

Diffuse growth and cytokinesis need microtubule nucleators, such as gamma tubulin and the gamma-tubulin ring complex [36]. Microtubules, in contrast to animal cells, may nucleate either from the nucleus, the plasma membrane, or from microtubules that were already there [34-37]. Gamma tubulin ring complexes are recruited during nucleation from preexisting microtubules by the two MOZART1 homolog proteins and AUGMIN complex, GAMMA-TUBULIN COMPLEX3 INTERACTING PROTEIN1 (GIP1) and GIP2 [38].

During the process of cytokinesis, the size of the phragmoplast increases as a result of the addition of additional microtubules that nucleate on the phragmoplast leading edge (Fig. 2). Nucleation that depends on microtubules, therefore, is essential for phragmoplast growth [39]. AUGMIN complexes are linked to microtubules during phragmoplast and spindle formation by ENDOSPERM DEFECTIVE1 (EDE1). Although the phragmoplasts of *ede1* mutants are deformed, there is no evidence of cytokinetic abnormalities [40]. Surprisingly, phragmoplast assembly does not need the GAMMA-TUBULIN COMPLEX PROTEIN6 (GCP6).

7. REGULATORY ROLE OF VARIOUS SEVERING PROTEINS DURING MICROTUBULE ARRAY REORIENTATION

It is necessary for Arabidopsis to have both the catalytic p60 KTN subunit and the microtubule cross-over targeting p80 KTN subunit in order to perform appropriate microtubule severing [41]. Microtubule reorientation in response to mechanical stress requires KTN1, which

suggests that severing helps microtubules realign [42]. Phragmoplast growth, location, and shape are all dependent on KTN1 [43]. Fragile fiber2 (*fra2*) mutants, also known as *ktn1*, exhibit decreased callose deposition and irregular cell plates [44]. At the distal phragmoplast, microtubules are severed by KTN1, which is hypothesized to be recruited by the microtubule-binding protein MACET4/CORD4 [22,35,45]. Though it has been shown that MACET4/CORD4 overexpression does not result in a rise in severing in tobacco (*Nicotiana tabacum*) cultivated cells, this has not been seen in tobacco (*Nicotiana tabacum*) cultured cells [46]. Rapid phragmoplast growth necessitates the presence of both KTN1 and MACET4/CORD4 [19-22,47].

Microtubule orientation and reconfiguration in response to mechanical signals are all impaired in the *ktn1* mutant cell [48]. Blue light-induced reorientation is impossible without KTN1 [26,49]. On the other hand, blue light-induced microtubule and mechanical force-dependent reorientation, are likely to have distinct processes.

8. SIGNIFICANCE OF MICROTUBULE BUNDLING PROTEINS DURING DIFFUSE GROWTH AND CYTOKINESIS

MOR1/GEM1 (Microtubule Organization1/Gemini1) is a critical component of microtubule assembly. Disruption of cytokinesis and diffuse growth occurs in *mor1* mutants [25,39,50]. The HEAT-repeat/TOG microtubule-binding domains of MOR1 are highly conserved [51]. Mitotic microtubule arrays, as well as cortical microtubules, include MOR1, which promotes the bundling of microtubules and their polymerization [52]. It has been shown that the temperature-sensitive *mor1* mutants may be exploited to better understand how microtubule dynamics and abnormal cell plate formation are regulated [53]. The phragmoplasts of *mor1/gem1* mutants occasionally split, resulting in many aberrant fresh cell walls and uneven membrane buildup in cell plates, in addition to generalized growth abnormalities [54]. It is interesting to note that despite the loss of parallel cortical microtubules and a reduction in the microtubule polymer mass in the *mor1* mutant at restricted temperature, cell wall crystallinity remained high. This suggests that there is an inverse relationship between microtubule mass and cellulose crystallinity [44,55].




Fig. 2. Model of land-plant cytokinesis. Phragmoplast, membrane, and cell wall structures that form the cell plate from left to right: vesicle accumulation, vesicle fusion to form a tubulo-vesicular network, a tubular network, and fenestrated sheet

MAP65-1 and MAP65-2 have functions in cytokinesis that are somewhat redundant with those of MAP65-3. This was discovered by studying the interactions of mutants with map65-3. Synthetic cytokinesis abnormalities are seen in the double mutants of map65-1 and map65-2 [56]. Although they generate fewer cells, map65-1 map65-2 double mutants do not show any evident cytokinesis deficiencies, indicating that they may still be active in cell division [49-51,57]. It is interesting to note that MAP65-1 may be seen at the very leading edge of the phragmoplast, where it crosslinks microtubules that run in the opposite direction [58]. Even though there is no abnormal phenotype in map65-4 mutants alone, when they are paired with map65-3 mutants, cytokinesis deficiencies induce gametophyte death [59]. The phragmoplast or phragmoplast midline is where MAP65-4 is found [32-35,60]. MAP65-4 is more dominant in the phragmoplast midline in the map65-3 mutant to partly substitute the antiparallel bundling role of MAP65-3 [61]. In vitro, MAP65-4 stimulates the formation of antiparallel and parallel microtubules [62]. Even though it may be found in the division area, the function of MAP65-4 is not understood at this time [63]. This significant family of microtubule-bundling proteins will benefit from a more combinatorial mutant study, which will reveal distinct and complementary roles.

9. KINESINS MEDIATED MOVEMENT OF CELLULAR COMPONENTS

Kinesins are microtubule and actin filament or microtubule-binding motor proteins that transport vesicles and organelles as well as actin filaments or microtubules. Kinins may be used to replace the uncommon minus-end directed dyneins found in plants [64]. It is thought that Golgi vesicles are delivered to the cell plate by plus-end directed kinesins that are located on the surface of the phragmoplast. In point of fact, even synthetic vesicles are able to be carried effectively to the cell plate that is in the process of development [65]. Despite the fact that electron microscopy has shown phragmoplast microtubules with associated vesicles, specific kinesins have yet to be found [66]. Kinesins, on the other hand, serve a variety of functions in cytokinesis and phragmoplast placement, showing the difficulty of coordinating cell plate development [61,64,67].

10. PLUS-END-DIRECTED KINESINS

Depending on the plant species, kinesins of the KINESIN-4 family have varied functions in cell growth and cytokinesis [22,34,56,68]. For the first time, an inflorescence stem with low mechanical strength was shown to be lacking in FRA1 (FRAGILE FIBER1) [49,55,69]. The plus-end

directed kinesin FRA1 is very processive [70]. However, further studies show that fra1-5 does not affect the organization, mobility, or concentration of CSCs at the plasma membrane and that the reduced mechanical strength in fra1 is not due to errors in cellulose microfibril patterning [71]. CMUs (cellulose synthase microtubule-uncoupling proteins) are proteins that bind microtubules and localize to the cell plate in order to prevent cells from twisting, and FRA1 does not directly alter cellulose production [72]. Microtubule lateral movement is controlled by modifying the FRA1-CMU association. The transport of polysaccharides to the apoplast may be affected by microtubule separation from the cortex, leading to fra1-cell wall abnormalities [73]. In point of fact, fra1-5 causes a reduction in the amount of fucose-alkyne-labeled pectin that is secreted [74]. In contrast, the cytokinesis of fra1-5 is unaffected [48,75]. During cytokinesis, mutants have a decreased number of cells but no visible structural problems in microtubules [15,65,76]. This protein family plays an important function in cytokinesis, however, in moss the Kin4-1a and Kin4-1c proteins are found in the phragmoplast midline, where they help to reduce microtubule overlapping areas [77]. Interesting, the phragmoplast of the kin4-1a and kin4-1c double mutant does not rotate, suggesting problems in the division plane [78].

Cytokinesis relies on a number of kinesin-7 plus-end directed kinesins. Tobacco possesses two kinesin-7s, NACK2 and NACK1, which have MSA (M-phase specific activator) sequences [79]. Mutations in the genes encoding tobacco HINKEL/NACK11 and Arabidopsis TETRASPORE / NACK2/STUD, as well as moss NACKa-c, were later shown to cause cytokinesis abnormalities [79,80]. Phragmoplast growth necessitates the microtubule-binding kinase-like protein RUNKEL, which is required for HINKEL recruitment to the phragmoplast [81]. Cell plate construction material may be transported by NACKs, since they are processive plus-end directed kinesins [81,82]. The C-terminal stalk domain of NACK1/HINKEL attracts MARKKK (mitogen-activated protein kinase kinase) proteins to the phragmoplast midline [83]. Proteases EB1, MAP65-1, and MAP65-3 are phosphorylated by the Mitogen-Activated Protein Kinase (MAPK) [84]. To ensure timely phragmoplast growth, Aurora Kinase activation of MAP65-1 lowers microtubule binding, which is important for appropriate microtubule disintegration at the phragmoplast trailing edge [85]. Extra Spindle Poles1 is a caspase-like

protein that stimulates microtubule polymerization with another set of kinesin-7s, which are not needed for cytokinesis [77,84,86].

11. MINUS END-DIRECTED KINESINS

In moss, Arabidopsis, and tobacco, members of the minus-end-directed Kinesin-5 family play an essential function in the process of cytokinesis; nevertheless, it is possible that they do not directly contribute to diffuse growth. Tobacco Kinesin-5, also known as Kinesin-Related Protein (NtKRP), is found in microtubules, where it is responsible for the translocation of opposite phragmoplast microtubules to reduce the amount of overlap between them [61,87]. Similar to Kinesin-5 AtKRP125c, rsw7 suffers spindle collapse, a faulty phragmoplast assembly and other problems in the cytokinesis process. It is possible that the apparent shape errors seen in cells are due to the mutant phenotypic changes that occur during mitosis and cytokinesis rather than flaws in swelling cells seen in RSW7 and AtKRP125c-deficient individuals [88]. Knocking down all Kinesin-5 (a-d) homolog RNAi has the same effect on the cell ability to divide into new cells. The minus-end-directed movement depletes Kinesin-5s from the phragmoplast midline, which is counterintuitive [89]. When Kinesin-5 is exempted from the deposition of antiparallel microtubules at the midline, it is not apparent how it connects or moves antiparallel microtubules.

During cytokinesis, a number of minus-end-directed kinesins from the Kinesin-14 family are involved. Spindle generation, chloroplast and nuclear mobility all rely on Kinesin-14s [57-60,90]. The vks1 mutant causes cytokinesis problems throughout endosperm cellularization, but not during vegetative phase, since it is essential for nuclear mobility, spindle formation, and phragmoplast organization [91]. Other similar proteins may be able to fill in during vegetative cell division, as this implies. For meiotic drive, a maize VKS1 paralog called KINDR moves neocentromeres toward the spindle poles; however, KINDR plays no function in vegetative cell division [91,92]. It is a calponin homology (KCH, KCH, calponin homology domains) kinesin, which attaches to both microtubules and actin and is found in mitotic microtubule frameworks [93].

12. CELLULOSE BIOSYNTHESIS

A substantial structural component of the cell wall is made up of cellulose. Long beta-1,4-linked

glucan chains make up the bulk of this substance. A huge complex called the CSC produces cellulose at the plasma membrane. Freeze fracture was used to create a hexameric rosette structure, which allowed researchers to see the CSC at the plasma membrane underneath microtubules [91,94]. CESA6 (CELLULOSE SYNTHASE 6), a CSC protein, was tracked in vivo using live imaging at the plasma membrane, where microtubules were shown to be a common route for its migration [78,95]. It is not known what the precise components of the CSC in primary cell walls are, however it is believed that they consist of a hexameric, equal-stoichiometric mixture of CESA1, CESA3, and either CESA6, CESA2, or CESA5 [84,96]. Cellular expansion and cytokinesis abnormalities may result from mutations in cellulose synthase subunits. Cytokinesis and cell elongation may be the sole roles for some of the *cesa6* mutants because of their redundant roles with CESA2 and CESA5 in the cell cycle [81,83,97].

Cell survival and movement may be improved by post-translational alterations such as phosphorylation [98]. Reduced mobility was seen in CESA5 and CESA3 and CESA3 by altering phosphorylation sites [99]. It is interesting that, phosphorylation of CESA by the protein kinase BRASSINOSTEROID INSENSITIVE2 is an intriguing example of the hormonal control of cellulose synthesis that occurs through phosphorylation of CESA [100].

13. PROTEINS ASSOCIATED WITH CELLULOSE SYNTHASE

A nebulous linker protein was long thought to be the mechanism by which the CSC connects to microtubules in the plasma membrane [97,101]. CSI1 (Cellulose Synthase Interactive 1), the first linker protein discovered, creates a physical interaction between CSCs and cortical microtubules to start and sustain structured and fast cellulose production (Fig. 3) [102]. Microtubule contact is confined to CSI1 N- and C-terminal areas, whereas the central domain of CESA interacts directly with CSI1 in several sites, including the N-, C-, and center regions of CSI1 [103]. A mutation in the *csi1* gene causes CESA particles to lose their ability to follow microtubules, resulting in decreased velocities [104].

It has been shown that two homologs, CSI3 and the less well-known CSI2, have varied degrees of expression in various tissues. Mutant plants that lack the *CSIA1* and the *CSIA3* genes have no apparent abnormalities, while those that lack both genes exhibit slower particle movement, a smaller hypocotyl, and less cellulose content [104,105]. CSI1 and its homologs may or may not be required for cytokinesis, although this remains an open question. To find out whether they are necessary for cytokinesis, it would be necessary to create a triple mutant.




Fig. 3. Model of trafficking of non-cellulosic polysaccharides and cellulose in the primary cell wall

COBRA (COB), called for the mutant roots snake-like appearance, and KOBITO (KOB), which means "little" in Japanese, are two proteins that are less well known yet crucial for cell growth and cellulose production [106]. Stunted development, decreased cellulose deposition, and damaged cell walls are all characteristics of kobito mutants [79,92,107]. There is strong evidence that the plasma membrane protein encoded by KOBITO is localized to this membrane. As a glycosyl-phospholipid-anchored protein, COB is a member of the COB family. In both the Golgi and the cell wall, COB accumulates [108]. Disorganized cellulose microfibrils and a decreased cellulose content are seen in cob mutants [109]. Increased grain yields and root hair formation are the results of roothairless3, a monocot-specific COB homolog in maize [110]. Rice homologs of COB, such as BRITTLE CULM1 (BC1) and BC1-Like4 (BC1L4), similarly stimulate cell growth [109,111]. Through an N-terminal domain, BC1 binds cellulose in vitro and localizes vesicles to the cell wall and to the plasma membrane [88,112]. By interacting directly with cellulose crystals, it is theorized that COB affects the synthesis of cellulose [103,113].

14. THE ASSEMBLY AND LOCALIZATION OF CSC

Electron microscopy revealed rosette CSCs in the Golgi, indicating that this is where they are assembled [114]. When cytokinesis occurred, immunogold labelling confirmed the Golgi deposition of CESA3 [115]. High molecular weight CSC complexes in stello1 stello2 mutants are decreased due of their involvement in the assembly of STELLO1 and STELLO1, two Golgi-localized proteins called after the Greek word stello meaning "to arrange in order."

It is hypothesized that endocytosis regulates the CSC, a large plasma membrane-localized complex [116]. When cells divide, clathrin-mediated endocytosis eliminates CESA particles from the plasma membrane and center of the cell plate, respectively [117]. Clathrin-mediated endocytosis and cytokinesis need DRP1A, a dynamin-like protein that binds to the light chain of clathrin [118]. Because of the deficiencies in rsw9/drps1a, cells are unable to extend and undergo cytokinesis properly, which leads to cellulose deficit [119]. OsDRP2B, a gene related to dynamin, was found to be mutated in rice at the brittle culm 3 mutant locus and is likely essential for clathrin-mediated endocytosis. OsCESA4 protein accumulation is reduced in

rice culm 3 mutants, but OsDRP2B overexpression increases OsCESA4 protein accumulation in plasma membrane [120]. Proof for endocytosis emerged from in vitro and yeast-two hybrid experiments that showed that CESA interacts with the medium subunit of ADAPTER PROTEIN COMPLEX 2 (AP2M) in clathrin-mediated endocytosis. CESA concentration rises in ap2m deleted mutants because of a disruption in clathrin-mediated endocytosis [121]. A clathrin-mediated endocytosis adaptor known as TRANSDUCIN/WD40-2 (TWD40-2) is a part of the TPLATE complex, which dates back millions of years [122]. In addition to regulating development and cytokinesis, the TPLATE complex also controls CESA endocytosis, as shown by a rise in plasma membrane CESA particle density in the absence of TPLATE [123]. TPLATE and MUNISCIN-LIKE (TML) interact with CESA6 and facilitate its internalization from the plasma membrane via the TPLATE Complex [124]. The recent development of a temperature-sensitive tplate mutant is going to be a very useful tool for elucidating the function that it plays in both diffuse proliferation and cytokinesis.

15. TRANSPORTATION AND BIOSYNTHESIS OF NON-CELLULOSIC POLYSACCHARIDE

Polysaccharides that are not cellulosic are mostly produced in the Golgi rather than at plasma membrane or late cell plate, as is the case with cellulose [125]. Xyloglucan and pectin are among the principal cell wall polysaccharides described here. Antibodies that have been well-studied and well-characterized may be a valuable tool in detecting certain carbohydrate characteristics [126]. Parallel to cellulose, xyloglucans have extra sugar alterations on the branching side chains of the glucans. According to the immunolocalization of terminal fucose antigens, modification of xyloglucans most likely takes place inside the trans cisternae of the Golgi and the TGN [127]. In agreement with this idea, oligosaccharide-mass characterization of Golgi-enriched subsets indicated fewer substituted xyloglucans in comparison to apoplastic cell wall segments [128]. Due to abnormally large secretory vesicles in phosphatidylinositol 4-kinase b1 and b2 (pi4kb1 and pi4kb2) mutants, xyloglucan trafficking from the trans Golgi to the TGN and to the cell surface is seen [129]. Golgi cis and medial cisternae are stained with immunogold, revealing the presence of xyloglucan-synthesizing enzymes such xylosyltransferases (XT1) and

galactosyltransferases MUR3 [130]. Overexpression of XT1 and MUR3 may lead to their being mislocalized in the cis and medial Golgi, but further research is needed to determine if xyloglucan side chain initiation occurs in early Golgi divisions.

Disorders in cell wall growth and the production of cytokinesis stubs have been found in mutants [131]. The cell plate accumulates CSLD5 in the form of punctate structures [132]. Mannans or xylans may be the cell wall substance that this enzyme generates [133]. However, the cytokinesis failure in Arabidopsis is exacerbated when the *csld5* mutant is co-dominant in combination with the *csld2* or *csld3* mutants, which normally govern root-hair development [134]. We may get vital insight into these proteins functions by clearly distinguishing which cell wall components they make.

16. CELL PLATE

An area without overlapping between microtubules and the cell plate assembly matrix is where vesicles arrive. This is where the cell plate assembly matrix may be found [135]. During cell plate development, monomeric RAB-GTPases link vesicles and facilitate vesicle transportation [136]. In order to go from the Golgi and TGN to the cell plate, RABGTPases such as RAB-E1, RAB-A3, and RAB-A2 are required [137]. RAB-A5c, which is a different kind of RAB-GTPase, tends to collect in its own distinct vesicles and rarely with the TGN. Cytokinesis and diffuse development are dependent on RAB-A5c, which is found on the cell plate [138]. Triple mutants of RAB-A1a-c are more susceptible to the cytokinesis-inducing drug Endosidin1 than single mutants [139]. STOMATAL CYTOKINESIS DEFECTIVE1 (SCD1), SCD2 and the Transport Protein Particle II (TRAPP II) complex components are also essential for cell plate formation [140]. Subsequently, the TRAPP-Interacting Plant Protein (TRIPP) was discovered to be a plant-specific component of the highly conserved complex of TRAPP. TGN to cell plate trafficking of TRIPP is essential for cytokinesis [141]. A fascinating interaction between exocyst proteins and SCD2, SCD1, and TRAPP II members may help coordinate vesicle trafficking at early and late stages during the development of cell plates [142].

17. CALLOSE: A MAJOR PRECURSOR OF CELL PLATE

Except in plasmodesmata and under circumstances that cause stress or injury,

callose, a significant component of the cell plate, is seldom seen in primary cell walls [143]. Xyloglucans and cellulose both include beta-1,4-linked glucan chains, but cellulose is made up of beta-1,3-linked glucans. When it comes to plasmodesmata, wound healing, pathogen response, pollen surface patterning, and other aspects of symplastic transport, many callose synthases play a role that is both distinct and overlapping [144]. Cellulosic mixtures with high levels of callose, which may be crucial for cell plate development, are more elastic in vitro [145]. Callose polymerization may provide the disseminating power needed to compress the tubulo-vesicular network into a fenestrated film, according to biophysical modelling.

Calcium is required for the action of callose synthase [146]. The ER may provide calcium to callose synthases during the development of cell plates: In several plant species, ER-cell plate connection has been observed [147]. Researchers have used staining to show that *Tradescantia virginiana* experiences calcium sequestration, which causes problems with cell plate development [148]. Mutants of the *P. patens* species have ER connections with the cell plate that are defective in the *sabre* strain, and this leads to abnormal callose buildup and ultimately cytokinesis abnormalities [149]. ER connections that have been changed may be responsible for the sluggish buildup of calloses in the *sabre* mutant. This is an intriguing possibility.

18. CONCLUSION

Modification and building of the cell wall are dynamic processes that are carefully regulated dynamic processes that include secretion and endocytosis. There are still many unanswered questions, such as how the addition of cellulose and non-cellulosic polymers is linked, how non-cellulosic polymers are transportation between the Golgi and the plasma membrane, and on which cytoskeletal tracks, and how the cytoskeleton and the activity of CSCs are significantly affected by environmental and mechanical gestures.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Abou-Saleh RH, Hernandez-Gomez MC, Amsbury S, Paniagua C, Bourdon M,

- Miyashima S, Helariutta Y, Fuller M, Budtova T, Connell SD, Ries ME. Interactions between callose and cellulose revealed through the analysis of biopolymer mixtures. *Nat. Commun.* 2018; 9(1):1-3.
2. Abu-Abied M, Belausov E, Hagay S, Peremyslov V, Dolja V, Sadot E. Myosin XI-K is involved in root organogenesis, polar auxin transport, and cell division. *J. Exp. Bot.* 2018;69(12):2869-2881.
 3. Ahn G, Kim H, Kim DH, Hanh H, Yoon Y, Singaram I, Wijesinghe KJ, Johnson KA, Zhuang X, Liang Z, Stahelin RV. SH3 domain-containing protein 2 plays a crucial role at the step of membrane tubulation during cell plate formation. *Plant Cell.* 2017;29(6):1388-1405.
 4. Ambrose C, Allard JF, Cytrynbaum EN, Wasteneys GO. A CLASP-modulated cell edge barrier mechanism drives cell-wide cortical microtubule organization in Arabidopsis. *Nat. Commun.* 2011;2(1):1-2.
 5. Ambrose C, Wasteneys GO. Microtubule initiation from the nuclear surface controls cortical microtubule growth polarity and orientation in Arabidopsis thaliana. *Plant Cell Physiol.* 2014;55(9):1636-1645.
 6. Ambrose JC, Shoji T, Kotzer AM, Pighin JA, Wasteneys GO. The Arabidopsis CLASP gene encodes a microtubule-associated protein involved in cell expansion and division. *Plant Cell.* 2007; 19(9):2763-2775.
 7. Amor Y, Haigler CH, Johnson S, Wainscott M, Delmer DP. A membrane-associated form of sucrose synthase and its potential role in synthesis of cellulose and callose in plants. *Proc. Natl. Acad. Sci. USA.* 1995; 92(20):9353-9357.
 8. Anderson CT. We be jammin: an update on pectin biosynthesis, trafficking and dynamics. *J. Exp. Bot.* 2016;67(2):495-502.
 9. Anderson CT, Carroll A, Akhmetova L, Somerville C. Real-time imaging of cellulose reorientation during cell wall expansion in Arabidopsis roots. *Plant Physiol.* 2010;152(2):787-796.
 10. Arieti RS, Staiger CJ. Auxin-induced actin cytoskeleton rearrangements require AUX1. *New Phytol.* 2020;226(2):441-459.
 11. Arima K, Tamaoki D, Mineyuki Y, Yasuhara H, Nakai T, Shimmen T, Yoshihisa T, Sonobe S. Displacement of the mitotic apparatuses by centrifugation reveals cortical actin organization during cytokinesis in cultured tobacco BY-2 cells. *J. Plant Res.* 2018;131(5):803-815.
 12. Asada T, Kuriyama R, Shibaoka H. TKRP125, a kinesin-related protein involved in the centrosome-independent organization of the cytokinetic apparatus in tobacco BY-2 cells. *J. Cell Sci.* 1997; 110(2):179-189.
 13. Assaad FF, Huet Y, Mayer U, Jürgens G. The cytokinesis gene KEULE encodes a Sec1 protein that binds the syntaxin KNOLLE. *J. Cell Biol.* 2001;152(3):531-544.
 14. Atmodjo MA, Sakuragi Y, Zhu X, Burrell AJ, Mohanty SS, Atwood III JA, Orlando R, Scheller HV, Mohnen D. Galacturonosyltransferase (GAUT) 1 and GAUT7 are the core of a plant cell wall pectin biosynthetic homogalacturonan: galacturonosyltransferase complex. *Proc. Natl. Acad. Sci. USA.* 2011;108(50):20225-20230.
 15. Bacete L, Melida H, Miedes E, Molina A. Plant cell wall-mediated immunity: cell wall changes trigger disease resistance responses. *Plant J.* 2018;93(4):614-636.
 16. Rehman RS, Zafar SA, Ali M, Pasha AN, Naveed MS, Waseem M, Raza A. CRISPR-Cas Mediated Genome Editing: A Paradigm Shift towards Sustainable Agriculture and Biotechnology. *Asian P. Res. J.* 2022;9(1):27-49.
 17. Bach L, Gissot L, Marion J, Tellier F, Moreau P, Satiat-Jeunemaître B, Palauqui JC, Napier JA, Faure JD. Very-long-chain fatty acids are required for cell plate formation during cytokinesis in Arabidopsis thaliana. *J. Cell Sci.* 2011;124(19):3223-3234.
 18. Baluška F, Liners F, Hlavačka A, Schlicht M, Van Cutsem P, McCurdy DW, Menzel D. Cell wall pectins and xyloglucans are internalized into dividing root cells and accumulate within cell plates during cytokinesis. *Protoplasma.* 2005;225(3): 141-155.
 19. Bannigan A, Scheible WR, Lukowitz W, Fagerstrom C, Wadsworth P, Somerville C, Baskin TI. A conserved role for kinesin-5 in plant mitosis. *J. Cell Sci.* 2007;120(16): 2819-2827.
 20. Baquero Forero A, Cvrčková F. SH3Ps— evolution and diversity of a family of proteins engaged in plant cytokinesis. *Int. J. Mol. Sci.* 2019;20(22):5623-5635.
 21. Bashline L, Li S, Anderson CT, Lei L, Gu Y. The endocytosis of cellulose synthase in

- Arabidopsis is dependent on $\mu 2$, a clathrin-mediated endocytosis adaptin. *Plant Physiol.* 2013;163(1):150-160.
22. Bashline L, Li S, Zhu X, Gu Y. The TWD40-2 protein and the AP2 complex cooperate in the clathrin-mediated endocytosis of cellulose synthase to regulate cellulose biosynthesis. *Proc. Natl. Acad. Sci. USA.* 2015;112(41):12870-12875.
 23. Baskin TI, Meekes HT, Liang BM, Sharp RE. Regulation of growth anisotropy in well-watered and water-stressed maize roots. II. Role of cortical microtubules and cellulose microfibrils. *Plant Physiol.* 1999; 119(2):681-692.
 24. Beeckman T, Przemeck GK, Stamatiou G, Lau R, Terryn N, De Rycke R, Inzé D, Berleth T. Genetic complexity of cellulose synthase A gene function in Arabidopsis embryogenesis. *Plant Physiol.* 2002; 130(4):1883-1893.
 25. Rehman RS, Zafar SA, Ali M, Ahmad M, Pasha AN, Waseem M, Hafeez AH, Raza A. Plant Pan-genomes: A New Frontier in Understanding Genomic Diversity in Plants. *J. Adv. Bio. Biotech.* 2022;25(1):10-22.
 26. Bernal AJ, Jensen JK, Harholt J, Sørensen S, Moller I, Blaukopf C, Johansen B, De Lotto R, Pauly M, Scheller HV, Willats WG. Disruption of ATCSLD5 results in reduced growth, reduced xylan and homogalacturonan synthase activity and altered xylan occurrence in Arabidopsis. *Plant J.* 2007;52(5):791-802.
 27. Bischoff V, Desprez T, Mouille G, Vernhettes S, Gonneau M, Höfte H. Phytochrome regulation of cellulose synthesis in Arabidopsis. *Curr. Biol.* 2011; 21(21):1822-1827.
 28. Rehman RS, Zafar SA, Ali M, Bxashir H, Pasha AN. Regulatory Role of Nucleus in miRNA Processing at Early Developmental Stages in Plants. *Asian J. Biotech. Gen. Engg.* 2022;5(4):8-11.
 29. Boruc J, Weimer AK, Stoppin-Mellet V, Mylle E, Kosetsu K, Cedeño C, Jaquinod M, Njo M, De Milde L, Tompa P, Gonzalez N. Phosphorylation of MAP65-1 by Arabidopsis Aurora kinases is required for efficient cell cycle progression. *Plant Physiol.* 2017;173(1):582-599.
 30. Bouquin T, Mattsson O, Næsted H, Foster R, Mundy J. The Arabidopsis lue1 mutant defines a katanin p60 ortholog involved in hormonal control of microtubule orientation during cell growth. *J. Cell Sci.* 2003; 116(5):791-801.
 31. Boutté Y, Frescatada-Rosa M, Men S, Chow CM, Ebine K, Gustavsson A, Johansson L, Ueda T, Moore I, Jürgens G, Grebe M. Endocytosis restricts Arabidopsis KNOLLE syntaxin to the cell division plane during late cytokinesis. *EMBO J.* 2010; 29(3):546-558.
 32. Boutté Y, Jonsson K, McFarlane HE, Johnson E, Gendre D, Swarup R, Friml J, Samuels L, Robert S, Bhalerao RP. ECHIDNA-mediated post-Golgi trafficking of auxin carriers for differential cell elongation. *Proc. Natl. Acad. Sci. USA.* 2013;110(40):16259-16264.
 33. Břejšková L, Hála M, Rawat A, Soukupová H, Cvrčková F, Charlot F, Nogué F, Haluška S, Žárský V. SEC6 exocyst subunit contributes to multiple steps of growth and development of *Physcomitrella* (*Physcomitrium patens*). *Plant J.* 2021; 106(3):831-843.
 34. Bringmann M, Li E, Sampathkumar A, Kocabek T, Hauser MT, Persson S. POM-POM2/cellulose synthase interacting1 is essential for the functional association of cellulose synthase and microtubules in Arabidopsis. *Plant Cell.* 2012;24(1):163-177.
 35. Brummell DA, CAMIRANDE A, Maclachlan GA. Differential distribution of xyloglucan glycosyl transferases in pea Golgi dictyosomes and secretory vesicles. *J. Cell Sci.* 1990;96(4):705-710.
 36. Burian A, Ludynia M, Uyttewaal M, Traas J, Boudaoud A, Hamant O, Kwiatkowska D. A correlative microscopy approach relates microtubule behaviour, local organ geometry, and cell growth at the Arabidopsis shoot apical meristem. *J. Exp. Bot.* 2013;64(18):5753-5767.
 37. Rehman RS, Pasha AN, Zafar SA, Ali M, Waseem M, Ahmad M, Ahmad N, Hafeez AH. Chromosomal Engineering through CRISPR/Cas Technology: A Way Forward. *J. Adv. Bio. Biotech.* 2022;25(1): 34-45.
 38. Rehman RS, Ali M, Zafar SA, Ahmad M, Pasha AN, Bashir H, Rashid F, Hussain M. Tapping into the Unsung Potential of CRISPR/CAS Technology in Agriculture. *Asian J. Biochem. Gen. Mol. Bio.* 2022; 10(4): 1-26.
 39. Burkart GM, Dixit R. Microtubule bundling by MAP65-1 protects against severing by inhibiting the binding of katanin. *Molecular*

- Biology of the Cell. 2019;30(13):1587-1597.
40. Buschmann H, Chan J, Sanchez-Pulido L, Andrade-Navarro MA, Doonan JH, Lloyd CW. Microtubule-associated AIR9 recognizes the cortical division site at preprophase and cell-plate insertion. *Curr. Biol.* 2006;16(19):1938-1943.
 41. Cao X, Deng M, Zhang ZL, Liu YJ, Yang XL, Zhou H, Zhang ZJ. Molecular characterization and expression analysis of TaTAC1 gene in *Triticum aestivum*. *J. Plant Genet. Resour.* 2017;18:125-132.
 42. Chen C, Zou J, Zhang S, Zaitlin D, Zhu L. Strigolactones are a new-defined class of plant hormones which inhibit shoot branching and mediate the interaction of plant-AM fungi and plant-parasitic weeds. *Sci. China C: Life Sci.* 2009;52(8):693-700.
 43. Chen Y, Dan Z, Gao F, Chen P, Fan F, Li S. Rice GROWTH-REGULATING FACTOR7 modulates plant architecture through regulating GA and indole-3-acetic acid metabolism. *Plant Physiol.* 2020; 184(1):393-406.
 44. Chen Y, Fan X, Song W, Zhang Y, Xu G. Over-expression of OsPIN2 leads to increased tiller numbers, angle and shorter plant height through suppression of OsLAZY1. *Plant biotechnol. J.* 2012;10(2): 139-149.
 45. Cui D, Neill SJ, Tang Z, Cai W. Gibberellin-regulated XET is differentially induced by auxin in rice leaf sheath bases during gravitropic bending. *J. Exp. Bot.* 2005; 56(415):1327-1334.
 46. Rehman RS, Ali M, Zafar SA, Bashir H, Pasha AN, Javed A, Ashraf NA, Shah WA. Regulatory Role of DNA Methylation and Its Significance in Plants. *Asian J. Biochem. Gen. Mol. Bio.* 2022;11(3):1-15.
 47. Dardick C, Callahan A, Horn R, Ruiz KB, Zhebentyayeva T, Hollender C, Whitaker M, Abbott A, Scorza R. PpTAC1 promotes the horizontal growth of branches in peach trees and is a member of a functionally conserved gene family found in diverse plants species. *Plant J.* 2013;75(4):618-630.
 48. Dayanandan P, Hebard FV, Baldwin VD, Kaufman PB. Structure of gravity-sensitive sheath and internodal pulvini in grass shoots. *Am. J. Bot.* 1977;64(10):1189-1199.
 49. Dharmasiri S, Swarup R, Mockaitis K, Dharmasiri N, Singh SK, Kowalchuk M, Marchant A, Mills S, Sandberg G, Bennett MJ, Estelle M. AXR4 is required for localization of the auxin influx facilitator AUX1. *Science.* 2006;312(5777):1218-1220.
 50. Dong H, Zhao H, Xie W, Han Z, Li G, Yao W, Bai X, Hu Y, Guo Z, Lu K, Yang L. A novel tiller angle gene, TAC3, together with TAC1 and D2 largely determine the natural variation of tiller angle in rice cultivars. *PLoS Genet.* 2016;12(11):112-120.
 51. Dong Z, Jiang C, Chen X, Zhang T, Ding L, Song W, Luo H, Lai J, Chen H, Liu R, Zhang X. Maize LAZY1 mediates shoot gravitropism and inflorescence development through regulating auxin transport, auxin signaling, and light response. *Plant Physiol.* 2013;163(3): 1306-1322.
 52. Edelmann HG. Graviperception in maize plants: is amyloplast sedimentation a red herring?. *Protoplasma.* 2018;255(6):1877-1881.
 53. Rehman RS, Zafar SA, Ali M, Pasha AN, Bashir H, Ashraf MA, Yaqoob MU, Hussain M. Biochemical and Molecular Insights into Abiotic Stress Tolerance in Plants. *Asian J. Biotech. Gen. Engg.* 2022;5(2):1-19.
 54. Fujihira K, Kurata T, Watahiki MK, Karahara I, Yamamoto KT. An agravitropic mutant of *Arabidopsis*, endodermal-amyloplast less 1, that lacks amyloplasts in hypocotyl endodermal cell layer. *Plant Cell Physiol.* 2000;41(11):1193-1199.
 55. Fukaki H, Fujisawa H, Tasaka M. Gravitropic response of inflorescence stems in *Arabidopsis thaliana*. *Plant Physiol.* 1996;110(3):933-943.
 56. Fukaki H, Wysocka-Diller J, Kato T, Fujisawa H, Benfey PN, Tasaka M. Genetic evidence that the endodermis is essential for shoot gravitropism in *Arabidopsis thaliana*. *Plant J.* 1998;14(4): 425-430.
 57. Gao H, Wang W, Wang Y, Liang Y. Molecular mechanisms underlying plant architecture and its environmental plasticity in rice. *Mol. Breed.* 2019;39(12):1-5.
 58. Rehman RS, Pasha AN, Zafar SA, Ali M, Bashir H, Saeed MU, Ashraf NA, Javed A. Molecular Mechanisms behind the Regulation of Rice Tiller Angle: An Update. *Asian J. Biol.* 2022;14(4):37-50.
 59. Geisler M, Blakeslee JJ, Bouchard R, Lee OR, Vincenzetti V, Bandyopadhyay A, Titapiwatanakun B, Peer WA, Bailly A, Richards EL, Ejendal KF. Cellular efflux of auxin catalyzed by the *Arabidopsis*

- MDR/PGP transporter AtPGP1. *Plant J.* 2005;44(2):179-194.
60. Gomez-Roldan V, Fermas S, Brewer PB, Puech-Pagès V, Dun EA, Pillot JP, Letisse F, Matusova R, Danoun S, Portais JC, Bouwmeester H. Strigolactone inhibition of shoot branching. *Nature.* 2008;455(7210):189-194.
 61. Harmoko R, Yoo JY, Ko KS, Ramasamy NK, Hwang BY, Lee EJ, Kim HS, Lee KJ, Oh DB, Kim DY, Lee S. N-glycan containing a core α 1, 3-fucose residue is required for basipetal auxin transport and gravitropic response in rice (*Oryza sativa*). *New Phytol.* 2016;212(1):108-122.
 62. Hashiguchi Y, Yano D, Nagafusa K, Kato T, Saito C, Uemura T, Ueda T, Nakano A, Tasaka M, Terao Morita M. A unique HEAT repeat-containing protein SHOOT GRAVITROPISM6 is involved in vacuolar membrane dynamics in gravity-sensing cells of Arabidopsis inflorescence stem. *Plant Cell Physiol.* 2014;55(4):811-822.
 63. Hollender C, Hill J, Waite J, Dardick C. Opposing influences of TAC1 and LAZY1 on lateral shoot orientation in Arabidopsis. *Sci. Rep.* 2020;10: 6051-6060.
 64. Rehman RS, Khan K, Zafar SA, Pasha AN, Ali M, Ashraf MA, Bashir H, Rashid F. Character Association Studies in Various Brassica napus Genotypes under Drought Stress. *Asian J. Res. Bot.* 2022; 7(2):20-27.
 65. Hollender CA, Waite JM, Tabb A, Raines D, Chinnithambi S, Dardick C. Alteration of TAC1 expression in Prunus species leads to pleiotropic shoot phenotypes. *Hortic. Res.* 2018;5:101-109.
 66. Hu M, Lv S, Wu W, Fu Y, Liu F, Wang B, Li W, Gu P, Cai H, Sun C, Zhu Z. The domestication of plant architecture in African rice. *Plant J.* 2018;94(4):661-669.
 67. Hu Y, Li S, Fan X, Song S, Zhou X, Weng X, Xiao J, Li X, Xiong L, You A, Xing Y. OsHOX1 and OsHOX28 redundantly shape rice tiller angle by reducing HSFA2D expression and auxin content. *Plant Physiol.* 2020;184(3):1424-1437.
 68. Huang L, Wang W, Zhang N, Cai Y, Liang Y, Meng X, Yuan Y, Li J, Wu D, Wang Y. LAZY2 controls rice tiller angle through regulating starch biosynthesis in gravity-sensing cells. *New Phytol.* 2021; 231(3):1073-1087.
 69. Rehman RS, Khan K, Zafar SA, Pasha AN, Ali M, Ashraf MA, Bashir H, Rashid F. Character Association Studies in Various Brassica napus Genotypes under Drought Stress. *Asian J. Res. Bot.* 2022;7(2):20-27.
 70. Huang X, Yang S, Gong J, Zhao Q, Feng Q, Zhan Q, Zhao Y, Li W, Cheng B, Xia J, Chen N. Genomic architecture of heterosis for yield traits in rice. *Nature.* 2016; 537(7622):629-633.
 71. Jiang J, Tan L, Zhu Z, Fu Y, Liu F, Cai H, Sun C. Molecular evolution of the TAC1 gene from rice (*Oryza sativa* L.). *J. Genet. Genomics.* 2012;39(10):551-560.
 72. Jin J, Huang W, Gao JP, Yang J, Shi M, Zhu MZ, Luo D, Lin HX. Genetic control of rice plant architecture under domestication. *Nat. Genet.* 2008;40(11):1365-1369.
 73. JONES JW, Adair CR. A "lazy" mutation in rice. *J. Hered.* 1938;29(8):315-318.
 74. Kaufman PB, Brock TG, Song IL, Rho YB, Ghosheh NS. How cereal grass shoots perceive and respond to gravity. *Am. J. Bot.* 1987;74(9):1446-1457.
 75. Rehman RS, Pasha AN, Zafar SA, Ali M, Bashir H, Saeed MU, Ashraf NA, Javed A. Molecular Mechanisms behind the Regulation of Rice Tiller Angle: An Update. *Asian J. Biol.* 2022;14(4):37-50.
 76. Kim JY, Ryu JY, Baek K, Park CM. High temperature attenuates the gravitropism of inflorescence stems by inducing SHOOT GRAVITROPISM 5 alternative splicing in Arabidopsis. *New Phytol.* 2016;209(1):265-279.
 77. Ku L, Wei X, Zhang S, Zhang J, Guo S, Chen Y. Cloning and characterization of a putative TAC1 ortholog associated with leaf angle in maize (*Zea mays* L.). *PLoS One.* 2011;6(6):123-130.
 78. Kunz HH, Häusler RE, Fettke J, Herbst K, Niewiadowski P, Gierth M, Bell K, Steup M, Flügge UI, Schneider A. The role of plastidial glucose-6-phosphate/phosphate translocators in vegetative tissues of Arabidopsis thaliana mutants impaired in starch biosynthesis. *Plant Biol.* 2010;12: 115-128.
 79. Lauri PE, Claverie J, Lespinasse JM. The effects of bending on the growth and fruit production of INRA Fercer® sweet cherry. *Acta. Hortic.* 1997:411-418.
 80. Ledesma N, Campbell RJ, Wasielewski J. Training and pruning a mango orchard to improve blooming and yield in South Florida. *Proc. Fla. State Hortic. Soc* 2016; 129:14-16.
 81. Li H, Sun H, Jiang J, Sun X, Tan L, Sun C. TAC4 controls tiller angle by regulating the endogenous auxin content and distribution

- in rice. *Plant Biotechnol. J.* 2021;19(1):64-73.
82. Rehman RS, Ali M, Zafar SA, Hussain M, Pasha A, Naveed MS, Ahmad M, Waseem M. Abscisic Acid Mediated Abiotic Stress Tolerance in Plants. *Asian J. Res. C. Sci.* 2022;7(1):1-17.
 83. Li Y, Deng Z, Kamisugi Y, Chen Z, Wang J, Han X, Wei Y, He H, Terzaghi W, Cove DJ, Cuming AC. A minus-end directed kinesin motor directs gravitropism in *Physcomitrella patens*. *Nat. Commun.* 2021;12(1):1-2.
 84. Li P, Wang Y, Qian Q, Fu Z, Wang M, Zeng D, Li B, Wang X, Li J. LAZY1 controls rice shoot gravitropism through regulating polar auxin transport. *Cell Res.* 2007;17(5):402-410.
 85. Li S, Zheng T, Zhuo X, Li Z, Li L, Li P, Qiu L, Pan H, Wang J, Cheng T, Zhang Q. Transcriptome profiles reveal that gibberellin-related genes regulate weeping traits in crape myrtle. *Hortic. Res.* 2020; 7-14.
 86. Li Y, Li J, Chen Z, Wei Y, Qi Y, Wu C. OsmiR167a-targeted auxin response factors modulate tiller angle via fine-tuning auxin distribution in rice. *Plant Biotechnol. J.* 2020;18(10):2015-2026.
 87. Li Y, Zhu J, Wu L, Shao Y, Wu Y, Mao C. Functional divergence of PIN1 paralogous genes in rice. *Plant Cell Physiol.* 2019; 60(12):2720-2732.
 88. Rehman RS, Zafar SA, Ali M, Pasha AN, Bashir H, Ashraf MA, Yaqoob MU, Hussain M. Biochemical and Molecular Insights into Abiotic Stress Tolerance in Plants. *Asian J. Biotech. Gen. Engg.* 2022;5(2):1-19.
 89. Li Z, Liang Y, Yuan Y, Wang L, Meng X, Xiong G, Zhou J, Cai Y, Han N, Hua L, Liu G. OsBRXL4 regulates shoot gravitropism and rice tiller angle through affecting LAZY1 nuclear localization. *Mol. Plant.* 2019;12(8):1143-1156.
 90. Liu JM, Mei Q, Xue CY, Wang ZY, Li DP, Zhang YX, Xuan YH. Mutation of G-protein γ subunit DEP1 increases planting density and resistance to sheath blight disease in rice. *Plant Biotechnol. J.* 2021;(3): 418-425.
 91. Marone D, Rodriguez M, Saia S, Papa R, Rau D, Pecorella I, Laidò G, Pecchioni N, Lafferty J, Rapp M, Longin FH. Genome-wide association mapping of prostrate/erect growth habit in winter durum wheat. *Int. J. Mol. Sci.* 2020;21(2): 394-400.
 92. Miyashita Y, Takasugi T, Ito Y. Identification and expression analysis of PIN genes in rice. *Plant Sci.* 2010; 178(5):424-428.
 93. Morita MT. Directional gravity sensing in gravitropism. *Annu. Rev. Plant Biol.* 2010; 61:705-720.
 94. Rehman RS, Ali M, Zafar SA, Bashir H, Pasha AN, Javed A, Ashraf NA, Shah WA. Regulatory Role of DNA Methylation and Its Significance in Plants. *Asian J. Biochem. Gen. Mol. Bio.* 2022;11(3):1-15.
 95. Morita MT, Tasaka M. Gravity sensing and signaling. *Curr. Opin. Plant Biol.* 2004;7(6): 712-718.
 96. Morita R, Sugino M, Hatanaka T, Misoo S, Fukayama H. CO₂-responsive CONSTANS, CONSTANS-like, and time of chlorophyll a/b binding protein Expression1 protein is a positive regulator of starch synthesis in vegetative organs of rice. *Plant Physiol.* 2015;167(4):1321-1331.
 97. Nagashima A, Uehara Y, Sakai T. The ABC subfamily B auxin transporter AtABC19 is involved in the inhibitory effects of N-1-naphthylphthalamic acid on the phototropic and gravitropic responses of *Arabidopsis* hypocotyls. *Plant Cell Physiol.* 2008;49(8):1250-1255.
 98. Najrana T, Sanchez-Esteban J. Mechanotransduction as an adaptation to gravity. *Front. Pediatr.* 2016;4:140-150.
 99. Nakamura M, Nishimura T, Morita MT. Gravity sensing and signal conversion in plant gravitropism. *J. Exp. Bot.* 2019; 70(14):3495-506.
 100. Rehman RS, Ali M, Zafar SA, Ahmad M, Pasha AN, Bashir H, Rashid F, Hussain M. Tapping into the Unsung Potential of CRISPR/CAS Technology in Agriculture. *Asian J. Biochem. Gen. Mol. Bio.* 2022; 10(4):1-26.
 101. Nakamura M, Toyota M, Tasaka M, Morita MT. An *Arabidopsis* E3 ligase, SHOOT GRAVITROPISM9, modulates the interaction between statoliths and F-actin in gravity sensing. *Plant Cell.* 2011;23(5): 1830-1848.
 102. Okamura M, Hirose T, Hashida Y, Ohsugi R, Aoki N. Suppression of starch synthesis in rice stems splay tiller angle due to gravitropic insensitivity but does not affect yield. *Funct. Plant Biol.* 2014;42(1):31-41.
 103. Okamura M, Hirose T, Hashida Y, Yamagishi T, Ohsugi R, Aoki N. Starch reduction in rice stems due to a lack of OsAGPL1 or OsAPL3 decreases grain

- yield under low irradiance during ripening and modifies plant architecture. *Funct. Plant Biol.* 2013;40(11):1137-1146.
104. You-nan O, Chun-sheng L, Shan-qing Z, Hui-min W, Lian-feng Z, Sheng-miao Y, Qian-yu J, Guo-ping Z. Dynamic changes of rice (*Oryza sativa* L.) tiller angle under effects of photoperiod and effective accumulated temperature. *Chin. J. Ecol.* 2009;20:56-65.
 105. Parker ML. Morphology and ultrastructure of the gravity-sensitive leaf sheath base of the grass *Echinochloa colonum* L. *Planta.* 1979;145(5):471-477.
 106. Periappuram C, Steinhauer L, Barton DL, Taylor DC, Chatson B, Zou J. The plastidic phosphoglucomutase from *Arabidopsis*. A reversible enzyme reaction with an important role in metabolic control. *Plant Physiol.* 2000;122(4):1193-1200.
 107. Rehman RS, Zafar SA, Ali M, Bashir H, Pasha AN. Regulatory Role of Nucleus in miRNA Processing at Early Developmental Stages in Plants. *Asian J. Biotech. Gen. Engg.* 2022;5(4):8-11.
 108. Rakusová H, Gallego-Bartolomé J, Vanstraelen M, Robert HS, Alabadí D, Blázquez MA, Benková E, Friml J. Polarization of PIN3-dependent auxin transport for hypocotyl gravitropic response in *Arabidopsis thaliana*. *Plant J.* 2011;67(5):817-826.
 109. Rijken PJ, De Groot RP, Kruijjer W, De Laat SW, Verkleij AJ, Boonstra J. Identification of specific gravity sensitive signal transduction pathways in human A431 carcinoma cells. *Adv. Space Res.* 1992;12(1):145-152.
 110. Roberts D. Identification of loci on chromosome 5A of wheat involved in control of cold hardiness, vernalization, leaf length, rosette growth habit, and height of hardened plants. *Genome.* 1990; 33(2):247-259.
 111. Sack FD. Plastids and gravitropic sensing. *Planta.* 1997;203(1):63-68.
 112. Rehman RS, Pasha AN, Zafar SA, Ali M, Waseem M, Ahmad M, Ahmad N, Hafeez AH. Chromosomal Engineering through CRISPR/Cas Technology: A Way Forward. *J. Adv. Bio. Biotech.* 2022;25(1):34-45.
 113. Sack FD, Leopold AC. Cytoplasmic streaming affects gravity-induced amyloplast sedimentation in maize coleoptiles. *Planta.* 1985;164(1):56-62.
 114. Sakuraba Y, Piao W, Lim JH, Han SH, Kim YS, An G, Paek NC. Rice ONAC106 inhibits leaf senescence and increases salt tolerance and tiller angle. *Plant Cell Physiol.* 2015;56(12):2325-2339.
 115. Sang D, Chen D, Liu G, Liang Y, Huang L, Meng X, Chu J, Sun X, Dong G, Yuan Y, Qian Q. Strigolactones regulate rice tiller angle by attenuating shoot gravitropism through inhibiting auxin biosynthesis. *Proc. Natl. Acad. Sci. USA.* 2014;111(30):11199-11204.
 116. Santelia D, Trost P, Sparla F. New insights into redox control of starch degradation. *Curr. Opin. Plant Biol.* 2015;25:1-9.
 117. Sieberer T, Leyser O. Auxin transport, but in which direction? *Science.* 2006; 312(5775):858-860.
 118. Slewinski TL, Braun DM. Current perspectives on the regulation of whole-plant carbohydrate partitioning. *Plant Sci.* 2010;178(4):341-349.
 119. Song K, Lee DW, Kim J, Kim J, Guim H, Kim K, Jeon JS, Choi G. EARLY STARVATION 1 is a functionally conserved protein promoting gravitropic responses in plants by forming starch granules. *Front. Plant Sci.* 2021;12-25.
 120. Rehman RS, Hussain M, Ali M, Zafar SA, Pasha AN, Bashir H, Ashraf NA, Javed A, Shah WA. A Comprehensive Review on Melatonin Compound and its Functions in Different Fungi and Plants. *Intern. J. Pathogen Res.* 2022;10(2):9-21.
 121. Strohm AK, Baldwin KL, Masson PH. Multiple roles for membrane-associated protein trafficking and signaling in gravitropism. *Front. Plant Sci.* 2012;3:274-285.
 122. Sun Q, Li TY, Li DD, Wang ZY, Li S, Li DP, Han X, Liu JM, Xuan YH. Overexpression of loose plant architecture 1 increases planting density and resistance to sheath blight disease via activation of PIN-FORMED 1a in rice. *Plant Biotechnol. J.* 2019;17(5):855-890.
 123. Swarup R, Kramer EM, Perry P, Knox K, Leyser HM, Haseloff J, Beemster GT, Bhalerao R, Bennett MJ. Root gravitropism requires lateral root cap and epidermal cells for transport and response to a mobile auxin signal. *Nat. Cell Biol.* 2005; 7(11):1057-1065.
 124. Tan L, Li X, Liu F, Sun X, Li C, Zhu Z, Fu Y, Cai H, Wang X, Xie D, Sun C. Control of a key transition from prostrate to erect growth in rice domestication. *Nat. Genet.* 2008;40(11):1360-1364.

125. Taniguchi M, Furutani M, Nishimura T, Nakamura M, Fushita T, Iijima K, Baba K, Tanaka H, Toyota M, Tasaka M, Morita MT. The Arabidopsis LAZY1 family plays a key role in gravity signaling within statocytes and in branch angle control of roots and shoots. *Plant Cell*. 2017;29(8):1984-1999.
126. Rehman RS, Zafar SA, Ali M, Ahmad M, Pasha AN, Waseem M, Hafeez AH, Raza A. Plant Pan-genomes: A New Frontier in Understanding Genomic Diversity in Plants. *J. Adv. Bio. Biotech*. 2022;25(1):10-22.
127. Tanimoto M, Tremblay R, Colasanti J. Altered gravitropic response, amyloplast sedimentation and circumnutation in the Arabidopsis shoot gravitropism 5 mutant are associated with reduced starch levels. *Plant Mol. Biol*. 2008;67(1):57-69.
128. Tian G, Gao L, Kong Y, Hu X, Xie K, Zhang R, Ling N, Shen Q, Guo S. Improving rice population productivity by reducing nitrogen rate and increasing plant density. *PLoS One*. 2017;12(8):99-109.
129. Tuncel A, Okita TW. Improving starch yield in cereals by over-expression of ADPglucose pyrophosphorylase: expectations and unanticipated outcomes. *Plant Sci*. 2013;211:52-60.
130. Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya Y, Shirasu K, Yoneyama K, Kozuka J. Inhibition of shoot branching by new terpenoid plant hormones. *Nature*. 2008;455(7210):195-200.
131. Overbeek JV. "LAZY," AN A-GEOTROPIC FORM OF MAIZE: "Gravitational Indifference" Rather Than Structural Weakness Accounts for Prostrate Growth-Habit of This Form. *J. Hered*. 1936;27(3):93-96.
132. Rehman RS, Hussain M, Ali M, Zafar SA, Pasha AN, Bashir H, Ashraf NA, Javed A, Shah WA. A Comprehensive Review on Melatonin Compound and its Functions in Different Fungi and Plants. *Intern. J. Pathogen Res*. 2022;10(2):9-21.
133. Vandenberg JP, Kiss JZ. Plant responses to gravity. *Semin. Cell Dev. Biol*. 2019;92:122-125.
134. Vitha S, Yang M, Sack FD, Kiss JZ. Gravitropism in the starch excess mutant of Arabidopsis thaliana. *Am. J. Bot*. 2007;94(4):590-598.
135. Vorselen D, Roos WH, MacKintosh FC, Wuite GJ, van Loon JJ. The role of the cytoskeleton in sensing changes in gravity by nonspecialized cells. *FASEB J*. 2014;28(2):536-547.
136. Waite JM, Dardick C. TILLER ANGLE CONTROL 1 modulates plant architecture in response to photosynthetic signals. *J. Exp. Bot*. 2018;69(20):4935-4944.
137. Rehman RS, Ali M, Zafar SA, Hussain M, Pasha A, Naveed MS, Ahmad M, Waseem M. Abscisic Acid Mediated Abiotic Stress Tolerance in Plants. *Asian J. Res. C. Sci*. 2022;7(1):1-17.
138. Wang JR, Hu H, Wang GH, Li J, Chen JY, Wu P. Expression of PIN genes in rice (*Oryza sativa* L.): tissue specificity and regulation by hormones. *Mol. Plant*. 2009;2(4):823-831.
139. Wang L, Xu Y, Zhang C, Ma Q, Joo SH, Kim SK, Xu Z, Chong K. OsLIC, a novel CCCH-type zinc finger protein with transcription activation, mediates rice architecture via brassinosteroids signaling. *PLoS One*. 2008;3(10):191-200.
140. Wang Y, Li J. Molecular basis of plant architecture. *Annu. Rev. Plant Biol*. 2008;59:253-279.
141. Liu J, Nannas NJ, Fu FF, Shi J, Aspinwall B, Parrott WA, Dawe RK. Genome-scale sequence disruption following biolistic transformation in rice and maize. *Plant Cell*. 2019;31(2):368-383.
142. Rehman RS, Zafar SA, Ali M, Pasha AN, Naveed MS, Waseem M, Raza A. CRISPR-Cas Mediated Genome Editing: A Paradigm Shift towards Sustainable Agriculture and Biotechnology. *Asian P. Res. J*. 2022;9(1):27-49.
143. Dawe RK. Charting the path to fully synthetic plant chromosomes. *Exp. Cell. Res*. 2020;390(1):111-126.
144. Coradini AL, Hull CB, Ehrenreich IM. Building genomes to understand biology. *Nat. Commun*. 2020;11(1):1-1.
145. Liu Y, Su H, Zhang J, Shi L, Liu Y, Zhang B, Bai H, Liang S, Gao Z, Birchler JA, Han F. Rapid birth or death of centromeres on fragmented chromosomes in maize. *Plant Cell*. 2020;32(10):3113-3123.
146. Mandáková T, Lysak MA. Post-polyploid diploidization and diversification through dysploid changes. *Curr. Opin. Plant Biol*. 2018;42:55-65.
147. Wolter F, Klemm J, Puchta H. Efficient in planta gene targeting in Arabidopsis using egg cell-specific expression of the Cas9

- nuclease of *Staphylococcus aureus*. Plant J. 2018;94(4):735-746.
148. Schindele P, Puchta H. Engineering CRISPR/LbCas12a for highly efficient, temperature-tolerant plant gene editing. Plant Biotechnol. J. 2020;18(5):1118.
149. Steinert J, Schiml S, Fauser F, Puchta H. Highly efficient heritable plant genome engineering using Cas9 orthologues from *Streptococcus thermophilus* and *Staphylococcus aureus*. Plant J. 2015; 84(6):1295-1305.

© 2022 Rehman et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91895>